

Reed Canary Grass (*Phalaris arundinacea L.*) – A multi-purpose crop June 2023

Authors: Dr Nana Afranaa Kwapong and Dr Elaine Jensen

Key messages:

- Reed canary grass (RCG) is a perennial, lignocellulosic crop that is native to the UK and has multiple uses.
- It is established from seed, making set-up costs lower than for many other perennial biomass crops.
- RCG is a shorter-term crop (5-10 years) than Miscanthus and willow, with average yields of 4-7 tonnes of dry matter per ha, depending on management.
- It is tolerant of a range of land types and conditions including flooding, drought, and freezing.
- RCG can be grazed in early summer (depending on the variety) and harvested for diverse other uses from late summer until early spring. This extends the annual



window of biomass availability and supports multiple end uses (e.g. anaerobic digestion, combustion, animal bedding, seed supply, game cover).

Biomass Connect is funded by the:



Introduction

Reed canary grass (*Phalaris arundinacea L.*, hereafter RCG) is a multipurpose, lignocellulosic perennial crop native to Europe, Asia, and North America. It can be grown on poorly productive land and tolerates a wide range of conditions. RCG is found in diverse habitats including wetlands and riparian zones but also drier areas like roadsides, forest margins, pastures, and disturbed areas. It is a fast-growing crop reaching heights of 1.5-2 m in a growing season, and, depending on management, can remain productive for 5-10 years.



Uses and benefits

- RCG can be <u>utilized in diverse energy conversion processes</u> like combustion, gasification, pyrolysis, and anaerobic digestion.
- RCG has a high gross calorific value (dry ~18.2 MJ kg⁻¹), exceeding that of wheat and Miscanthus. Ash content (~3-8 %) depends particularly on harvest, management, and site.
- RCG has carbon sequestration potential and can be used for a prolonged 'fallow' period in a rotation where soil health can recover from more intensive arable practices and organic matter can be accrued.
- RCG creates habitat for diverse wildlife including birds and small mammals.

Biomass Connect is funded by the:



- RCG can mitigate <u>greenhouse gas emissions</u> and reduce nitrate leaching, acting as a buffer crop.
- RCG can be integrated into <u>multifunctional landscapes</u> as a biomass crop providing benefits of supporting ecosystem services and mitigating environmental impacts.
- RCG can be used for livestock feeding provided it is low in alkaloids (eg. Marathon or Palaton variety). It has good absorption properties and is a viable alternative to straw as <u>bedding</u> <u>material for livestock</u>.
- RCG has potential to remove <u>contaminants</u> from soil and water.

Establishment

RCG is planted by seed. It requires a firm, smooth seedbed and timely weed control during establishment, with fertilizer where nitrogen is limiting. Seed can be broadcasted or planted using a conventional seed drill. Conventional seed drilling provides more uniform stand development and seed depth. The ideal time to plant RCG is between March and April when soils are moist, and the root system has plenty of time to establish. The recommended seeding rate for forage production is 5-10 kg ha⁻¹ with row spacing of 12.5 cm, at a depth of about 2 cm. Rolling is recommended both before and after sowing to conserve the moisture in the seedbed followed by application of broadleaf herbicide to control weeds. Growth can be slow during the first year of cultivation while the root and rhizome system establishes. Harvesting can usually commence from year two. Nutrients translocate to the rhizomes over the autumn and winter period, providing energy for new shoots in spring, and improving the sustainability of the crop and quality of the biomass for combustion purposes.

Site suitability

RCG is highly tolerant to waterlogging (established stands can tolerate extended periods of inundation) and a wide pH range from pH 4.9 to pH 8.2. It can thrive on marginal land unsuitable for food or feed production. Once established, Reed canary grass is one of the most productive temperate grasses during drought due to its deep rhizome and root system. As a temperate grass used extensively in Scandinavia, it is also highly tolerant of freezing conditions.

Biomass Connect is funded by the:



Biomass production and harvesting

Average yields of 5-7 tonnes per hectare are achievable in the UK, depending on management, although 'pushing' the crop may decrease crop overall persistence. Yields will be improved by fertilization where nutrients are limiting although over fertilizing may promote lodging. Harvesting the crop green will reduce sustainability, as nutrients not yet translocated to the below-ground biomass will be removed.



Reed canary grass under two harvesting regimes (autumn, LHS; spring RHS) on shallow, stony soil in West Wales.

Harvesting can be conducted with conventional grass harvesting machinery and usually commences from the second year after sowing. The timing of harvest depends on the intended market/end-use. Forage quality is best in spring and early summer, while harvest for anaerobic digestion can extend from summer until autumn senescence begins. For combustion and animal bedding, harvesting is best delayed until late winter or early spring, where combustion quality will be improved through nutrient translocation below ground, leading to lower contents of sodium, potassium, chlorine and moisture (ca. 10-15%). Spring-harvested biomass requires little or no drying. The crop is typically mown first before being baled and transported. Alternatively, the grass can be cut and chipped using a forage harvester. Eliminating the crop after productive years will be easiest using herbicides e.g., glyphosate, although repeated cutting and conventional tillage may suffice.

Harvesting RCG whilst green will remove nutrients, increasing the need for fertilizer, but research has shown costs can be reduced using <u>sewage sludge</u> or <u>nitrogen-fixing legumes</u>.

Biomass Connect is funded by the:



Pest and Diseases

RCG disease levels are relatively low and are not considered a cause for concern (they include brown rust, mildew, buff spot, powdery mildew, and Rhynchosporium (leaf scald)). However, RCG can be attacked by insect larvae, which feed inside stem bases and can occasionally significantly reduce yield. Damage from rabbits and slugs can occur during the first year of establishment.

Economics

Returns during the first year are low, as establishment costs will not be balanced by revenue from biomass production. A <u>study</u> conducted in the UK found the average gross margin of £153.89 ha⁻¹ for RCG grown in multiple sites. The gross margins ranged from a minimum of £110.60 ha⁻¹ to a maximum of £219.40 ha⁻¹ across the sites studied. The differences in the gross margin were attributed to the variations in yield among different sites and prices of the biomass.